REFERENCES

1. Alkhalidi BA, Tamimi JJ, Salem II, Ibrahim H and Sallam AA (2008). Assessment of the bioequivalence of two formulations of clarithromycin extended-release 500- mg tablets under fasting and fed conditions: a single dose, randomized, open-label, two-period, two-way crossover study in healthy Jordanian male volunteers. Clin. Ther., 30(10): 1831-1843.
2. Banka NH (2003). Role of prokinetics in dyspepsia. Gastroenterol Today, 7: 1-4.
3. Berkow R (1992). The Merck Manual, Merck & Co., Inc., Rahway, N.J., USA, 16th Ed, 42- 45.
4. Booth S L, Tucker K L, McKeown N M, Davidson K W, Dallal G E and Sadowski J A (1997). Relationship between dietary intakes and fasting plasma concentrations of fat-soluble vitamins in humans. J. Nutr., 127: 587– 592.
5. Charman WN, Porter CJ, Mithani S and Dressman JB (1997). Physio - chemical and physiological mechanisms for the effects of food on drug absorption: the role of lipids and pH. J. Pharm. Sci. 86: 269–282.
6. D' Arcy PF (1985). Nitrofurantoin. Drug Intell. Clin. Pharm., 19(7-8): 540-547.
7. Fassbender M, Lode H, Schaberg T, Borner K and Koeppe P (1993). Pharmacokinetics of new oral cephalosporins, including a new carbacephem. Clin. Infect. Dis., 16(5): 646-653.
8. Foulds G, Luke DR, Teng R, Willavize SA, Friedman H and Curatolo WJ (1996). The absence of an effect of food on the bioavailability of azithromycin administered as tablets, sachet or suspension. J. Antimicrob. Chemother., 37, 37-44.
9. Gibaldi M. (1991). Compartmental and noncompartmental pharmacokinetics. In Biopharmaceutics and Clinical Pharmacokinetics, 4: 14-23.
10. Hepler CD and Strand LM (1990). Opportunities and responsibilities in pharmaceutical care. Am. J. Hosp. Pharm., 47(3): 533-43.
11. Hokuriko Seiyaku Co. Ltd. Data on file. Revised: April 2001.
12. Hyun Chul Lim, Young Gyun Kim, Jung Hyun Lim, Hee Sun Kim, and Hyojin Park (2008). Effect of Itopride Hydrochloride on the Ileal and Colonic Motility in Guinea Pig In Vitr. Yonsei Med J, 49(3):472 – 478.

13. Inoue K, Sanada Y, Fujimura J and Mihara O (1999). Effect of Itopride hydrochloride on the digestive symptoms of chronic gastritis with reflux esophagitis. Clin Med, 15: 1803-09.
14. Iwanga Y, Kemura T, Miyashita N et al (1994). Characterisation of acetylcholinesterase inhibition by Itopride, Jpn J Pharmacol, 66: 317-22.
15. Iwanga Y, Miyashita N, MizutaniF, (1991). Stimulatory effect of N-[-4[2-(dimethyl-amino)ethoxy]benzyl]-3-4-dimethoxybenzamide hydrochloride (HSR-803) on normal and delayed gastrointestinal propulsion, Jpn J Pharmacol, 56: 261- 69.
16. Iwanga Y, Miyashita N, Morikawa K, Mizumoto A, Kondo Y, Itoh Z (1990). A novel water soluble dopamine-2 antagonist with anticholinesterase activity in gastrointestinal motor activity. Gastroenterol, 99: 57-64.
17. Iwanga Y, Suzuki N, Kato KI, Morikawa K, Kato H, Ito Y, Gomi Y (1993). Stimulatory effects of HSR-803 on ileal motor activity, Jpn J Pharmacol, 62: 395-401.
18. Johnson RD, Dorr MB, Hunt TL, Jensen BK and Talbot GH (1999). Effects of food on the pharmacokinetics of sparfloxacin. Clin. Ther., 21(6): 982-991.
19. Kakuichi M, Saito T and Ohara N (1997). Pharmacological evaluation of Itopride hydrochloride with regard to drug induced arrhythmia. Jpn Pharmacol Ther, 25: 822-27.
20. Kamath, Vinod K, Verghese J and Bhatia S (2003). Comparative evaluation of the efficacy and tolerability of Itopride and Metoclopramide in patients with NUD. JAMA, 2(8), 95-98.
21. Karim S, Ahmed T, Monif T, Saha N and Sharma PL (2003). The effect of four different types of food on the bioavailability of cefaclor. Eur. J. Drug Metab. Pharmacokinet., 28(3): 185-190.
22. Kawakami J, Matsuse T, Kotaki H, Seino T, Fukuchi Y, Orimo H, Sawada Y and Iga T (1994). The effect of food on the interaction of ofloxacin with sucralfate in healthy volunteers. Eur. J. Clin. Pharmacol., 47(1): 67- 69.
23. Kshirsagar NA and Ankalesaria PS (1987). Effect of food on doxycycline absorption. J. Postgrad. Med., 33:117- 119.
24. Knoben JE and Anderson PO (1993), Handbook of Clinical Drug Dam, Drug Intelligence Publications Inc., Hamilton Illinois, USA, 7: 76-78.

25. L. D. Bornemann, S. Castellano, A. H. Lin, D. Enthoven, and I. H. Patel, (1988). Influence of food on bioavailability of amdinocillin pivoxil, antimicrobial agents and chemotherapy, 32 (4): 592-594.
26. Lecaillon JB, Hirtz JL, Schoeller JP, Humbert G and Vischer W (1980). Pharmacokinetic comparison of cefroxadin (CGP 9000) and cephalexin by simultaneous administration to humans. Antimicrob. Agents Chemother., 18(5): 656-660.
27. Lehto P and Kivistö KT (1995). Effects of milk and food on the absorption of enoxacin. Br. J. Clin. Pharmacol., 39(2): 194-196.
28. Lee LJ, Hafkin B, Lee ID, Hoh J and Dix R (1997). Effects of food and sucralfate on a single oral dose of 500 milligrams of levofloxacin in healthy subjects. Antimicrob. Agents Chemother., 41(10): 2196-2200.
29. Marathe PH, Arnold ME, Meeker J, Greene DS and Barbhaiya RH (2000). Pharmacokinetics and bioavailability of a metformin/glyburide tablet administered alone and with food. J. Clin. Pharmacol., 40(12 Pt 2): 1494-1502.
30. Mantyla R, Männistö PT, Vuorela A, Sundberg S and Ottoila P (1984). Impairment of captopril bioavailability by concomitant food and antacid intake. Int. J. Clin. Pharmacol. Ther. Toxicol., 22(11): 626-629.
31. Marino MR and Vachharajani NN (2001). Drug interactions with irbesartan. Clin. Pharmacokinet., 40(8): 605-614.
32. Mcquaid KR and Mark Feldman (2002). Sleisenger&Fordtran's Dyspepsia In: Gastrointestinal and Liver Disease. Pathophysiology/Diagnosis/Management, 7(1): 102-18.
33. Micromedex Inc. 93: l974-l997.
34. Miyoshi A, Masamime O and Sekiguchi T (1994). Clinical evaluation of Itopride hydrochloride for gastrointestinal symptoms associated with chronic gastritis: a multicentre double blind clinical trial using cisapride as control drug. Clin Pharmacol & Therap, 4 (2): 261-79.
35. Mohammad Yaheya Mohammad Ismail (2009). Herb-drug interactions and patient counseling. International journal of pharmacy and pharmaceutical sciences, 1 (1): 151-161.

36. Mushiroda T, Douya R, Takahara E and Nagata O (2000). The involvement of flavin containing monooxygenase but not CYP3A4 in metabolism of Itopride hydrochloride, a gastrokinetic agent: comparison with cisapride and mosapride citrate. Drug Metab Dispos, 28: 1231-37.
37. Nakajima M, Uematsu T, Nakajima S, Nagata O and Yamaguchi T (1993). Phase 1 study of HSR-803. JpnPharmacolTher, 21(11): 4157-73z.
38. Nai-Ning Songa,b, Shao-Yu Zhangb and Chang-Xiao Liu (2004). Overview of factors affecting oral drug absorption. Asian Journal of Drug Metabolism and Pharmacokinetics, 4(3): 167-176.
39. Noda T (2001). Effects of meal on Pharmacokinetics of HSR-803 in Humans. Hokuriku Seiyaku Co Ltd. Data on file.
40. Noritake M, Kikuchy Y and Otsubo T (1997). Effect of Itopride hydrochloride on diabetic Gastroparesis. KisoToRinsho, 31(8): 2785-91.
41. Ohki R, Takahashi M, Mizuno O and Fujikawa H (2001). Torsades de pointes ventricular tachycardia induced by mosapride and flecainide in the presence of hypokalemia. Pacing ClinElectophysiol, 24: 119-21.
42. Otsuba T, Mizokami Y, Shiraishi T, Narasaka T, Nakamura H and Takeyama H (1998). Effect of Itopride hydrochloride on nonulcer dyspepsia. Clin Med, 14: 94-97.
43. Pargal A, Kelkar MG and Nayak PJ (1996). The effect of food on the bioavailability of ibuprofen and flurbiprofen from sustained release formulations. Biopharm. Drug Dispos., 17(6): 511-519.
44. Pasricha PJ. Prokinetic agents, antiemetics agents used in irritable bowel syndrome. In: Hardman JG (2001). Goodman and Gilman'ssThe Pharmacological Basis of Therapeutics, New York, McGraw Hill Book Inc., 10: 1021.
45. Santosh U Zate, Paresh I Kothawade, Mohan N Rathi, Mohan H Shitole, Chetan P Yewale and Vinayak S Gawande (2010). Development and characterization of gastroretentive mucoadhesive tablets of venlafaxine hydrochloride. International Journal of Drug Delivery, 2: 299-303.
46. Sawant P, Kalokhe S and Patil S (2002). Comparative evaluation of Itopride hydrochloride and domperidone in patients with dyspepsia and chronic gastritis. Gastroenterol Today, 2: 1-4.

47. Seema Gupta, V Kapoor and B Kapoor (2004). Itopride: A Novel Prokinetic Agent. JK Science, 6 (2): 106- 108.
48. Shah A, Liu MC, Vaughan D and Heller AH (1999). Oral bioequivalence of three ciprofloxacin formulations following single-dose administration: 500 mg tablet compared with 500 mg/10 mL or 500 mg/5 mL suspension and the effect of food on the absorption of ciprofloxacin oral suspension. J. Antimicrob. Chemother., 43(Suppl A): 49-54.
49. Shimomura K (1992).A dose finding experiment for a study by oral administration of HSR-803 during the period of fetal organogenesis in rabbits, CLEA Preclin Res Rep,18 (2): 145.
50. Song JC and White CM (2002). Clinical pharmacokinetics and selective pharmacodynamics of new angiotensin converting enzyme inhibitors: An update. Clin. Pharmacokinet., 41(3): 207-224.
51. Stocklcy IH (1996). Drug Interactions, The Pharmaceutical Press, London, UK, 4: 253-257.
52. Syed Faisal Aman, Fouzia Hassan, Baqar s Naqvi and Syed Muhammmad Farid Hasan (2010). Studies of food drug interactions, Pak. J. Pharm. Sci., 23(3):313-320.
53. Tadashi Tsubouchi, Takaharu Saito, FujieMizutani, ToshieYamauchi and YujiIwanga (2003). Stimulatory action of Itopride hydrochloride on colonic motor activity in vitro and in vivo. J Pharmacol Exp Therapeut, 306: 787-93.
54. Takuma K, Ohtani K, Kotaki H and Iga T (1997). Comparative studies of drug induced arrhythmia in guinea pigs by cisapride and Itopride hydrochloride: prolongation of QT interval and search for alternative drugs to avoid side effect. The Annual meeting of Hospital and Pharmaceutical society of Japan, Nagoya, 13-14.
55. The pharmaceutical codex: Principles and practice of pharmaceutics (1994). The Pharmaceutical Press, London, UK, 12: 536- 540.
56. Thomson AB, Sinclair P, Matisko A, Rosen E, Andersson T and Olofsson B (1997). Influence of food on the bioavailability of an enteric-coated tablet formulation of omeprazole 20 mg under repeated dose conditions. Can. J. Gastroenterol., 11(8):663-667.
57. Vasu S, Adithan C, Shashindran CH, Asad M, Koumaravelou K and Topno I (2000). Effect of two types of Indian breakfast on bioavailability of cefuroxime axetil. Indian J. Med. Res., 112: 104-108.

58. Williams L, Hill DP, Davis JA and Lowenthal DT (1996). The influence of food on the absorption and metabolism of drugs: an update. Eur J Drug Metab Pharmacokinet, 21:201-11.